chapter 9 linear momentum and collisions Full Version

chapter 9 linear momentum and collisions is a fundamental topic in classical physics that explores how objects move and interact during collisions. This chapter builds on the concepts of force and motion introduced earlier, focusing on the conservation laws that govern the behavior of objects when they collide. Understanding linear momentum and collisions is essential for analyzing a wide range of physical phenomena, from car crashes to atomic interactions, and it provides the foundation for more advanced topics in physics and engineering.

Understanding Linear Momentum

Definition of Linear Momentum

Linear momentum, often simply called momentum, is a vector quantity defined as the product of an object?s mass and its velocity:

- **Mathematically:** p = m × v
- **p** represents the momentum vector, which has both magnitude and direction.

This means that an object with a larger mass or a higher velocity has greater momentum.

The Significance of Momentum

Momentum is a crucial concept because it remains conserved in isolated systems. This conservation law allows physicists to predict the outcome of interactions without detailed knowledge of the forces involved. It also helps in understanding how energy and motion are transferred during collisions.

Conservation of Linear Momentum

Law of Conservation of Momentum

The principle states that in the absence of external forces, the total linear momentum of a system remains constant. Mathematically:

- **Initial total momentum = Final total momentum**

This principle is a cornerstone of classical mechanics and is valid for both elastic and inelastic collisions.

Implications of Conservation

The conservation law simplifies the analysis of collisions:

- It allows calculation of unknown velocities after a collision.
- It explains phenomena like recoil and transfer of motion.

When external forces act, they change the total momentum, but in an isolated system, this law holds true.

Types of Collisions

Elastic Collisions

In elastic collisions:

- Both kinetic energy and momentum are conserved.
- Objects bounce off each other without permanent deformation or energy loss.
- Examples include collisions between billiard balls or gas particles.

The mathematical conditions for an elastic collision involve both conservation of momentum and kinetic energy.

Inelastic Collisions

In inelastic collisions:

- Momentum is conserved, but kinetic energy is not.
- Some energy is transformed into heat, sound, or deformation.
- Perfectly inelastic collisions are a special case where objects stick together after collision.

Understanding the difference is essential for analyzing real-world interactions.

Mathematical Analysis of Collisions

One-Dimensional Collisions

In many problems, collisions occur along a single line, simplifying analysis:

- Identify initial velocities and masses of the objects.
- Apply conservation of momentum:

- m?v?i + m?v?i = m?v?f + m?v?f

- For elastic collisions, also apply conservation of kinetic energy:

- 0.5 m?v?i² + 0.5 m?v?i² = 0.5 m?v?f² + 0.5 m?v?f²

- Solve for the unknown final velocities.

Two-Dimensional Collisions

When collisions occur in a plane:

- Separate the velocities into components along axes (usually x and y).
- Apply conservation of momentum separately along each axis.
- Use vector addition to find the resultant velocities after collision.

Vector diagrams are helpful visual tools for understanding these interactions.

Impulse and Its Role in Collisions

Definition of Impulse

Impulse is the product of the force applied and the time interval during which it acts:

- **J = F × ?t**

It is a vector quantity and relates directly to the change in momentum:

- **J = ?p = p_final - p_initial**

Impulsive Forces

During a collision, forces act over a short time, producing an impulse:

- Impulses can be large even if forces are brief.
- They cause significant changes in momentum.

Understanding impulse helps in designing safety features like airbags and crumple zones in vehicles.

Real-World Applications of Momentum and Collisions

Automobile Safety and Crash Analysis

Engineers analyze collisions using momentum principles:

- Calculating impact forces.
- Designing safety features to minimize injury.
- Determining energy absorption during crashes.

Sports and Recreation

In sports:

- Ball trajectories after collisions with bats or rackets are predicted using momentum conservation.
- Understanding how players transfer momentum to objects enhances performance and safety.

Atomic and Subatomic Collisions

In particle physics:

- Collisions between particles reveal fundamental forces and particles.
- Accelerator experiments rely heavily on momentum conservation to analyze outcomes.

Summary and Key Takeaways

To summarize:

- Linear momentum is a vector quantity crucial for analyzing motion in collisions.
- The law of conservation of momentum states that total momentum remains constant in isolated systems.
- Collisions are classified as elastic or inelastic based on energy conservation.
- Impulse relates the force applied during collision to the change in momentum.
- Understanding these concepts is vital across various fields, from engineering to fundamental physics.

Further Study and Practice Problems

To deepen your understanding:

- Solve problems involving collisions between different masses moving at various initial velocities.
- Explore the effects of external forces on momentum conservation in real-world scenarios.
- Investigate the role of energy transformation in inelastic collisions.

Applying these principles through practical problems enhances comprehension and prepares you for more advanced physics topics.

chapter 9 linear momentum and collisions forms a cornerstone of physics education, offering insights into how objects interact and transfer motion. Mastery of these concepts enables students and professionals alike to analyze and predict the outcomes of complex physical interactions, fostering a deeper appreciation of the fundamental laws that govern our universe.

Understanding Chapter 9: Linear Momentum and Collisions is fundamental to grasping the principles governing the motion and interaction of objects in physics. Whether you're a student preparing for exams, an educator designing curriculum, or an enthusiast seeking to deepen your knowledge, this comprehensive guide will walk you through the core concepts, mathematical foundations, and real-world applications of linear momentum and collisions.

Introduction to Linear Momentum

Linear momentum is a vector quantity that describes the motion of an object and is defined as the product of its mass and velocity. It encapsulates the idea of an object's motion in a way that remains consistent across different frames of reference. In simple terms, linear momentum tells us how difficult it is to change an object's motion.

Key Definition:

- Linear Momentum (p) = Mass (m) × Velocity (v)

This fundamental quantity is conserved in isolated systems, forming the backbone of collision analysis.

Fundamental Principles of Linear Momentum

Conservation of Momentum

One of the most important principles in physics is the law of conservation of linear momentum. It states that in an isolated system?meaning no external forces act on the objects involved?the total momentum before an interaction equals the total momentum after.

Mathematically:

- ? p_initial = ? p_final

This principle is essential when analyzing collisions and explosions because it allows us to predict the outcome of interactions without knowing the detailed internal forces.

Implication in Collisions

In collisions, momentum conservation helps determine unknown velocities after impact, especially when combined with other conservation laws such as energy conservation (though energy might not always be conserved in inelastic collisions).

Types of Collisions

Collisions are categorized based on whether kinetic energy is conserved:

Elastic Collisions

- Both momentum and kinetic energy are conserved.
- Occur in idealized situations; for example, collisions between billiard balls or gas particles.
- After collision, objects bounce off each other without permanent deformation or heat generation.

Inelastic Collisions

- Momentum is conserved, but kinetic energy is not.
- Some energy is transformed into internal energy, heat, or sound.
- Real-world collisions are often inelastic; for example, car crashes.

Perfectly Inelastic Collisions

- The colliding objects stick together after impact.
- The maximum possible loss of kinetic energy for a given momentum exchange.
- Example: a lump of clay hitting and sticking to a surface.

Mathematical Treatment of Collisions

Understanding the equations governing collisions is key to solving physics problems involving momentum.

Conservation of Momentum Equation

For two objects with masses m_1 and m_2, initial velocities v_{1i} and v_{2i}, and final velocities v_{1f} and v_{2f}:

$$m_1 v_{1i} + m_2 v_{2i} = m_1 v_{1f} + m_2 v_{2f}$$

This equation applies to all types of collisions, but additional considerations are needed for energy in elastic collisions.

Elastic Collisions: Using Both Momentum and Kinetic Energy

In elastic collisions, both momentum and kinetic energy are conserved:

- Momentum conservation:

$$m_1 v_{1i} + m_2 v_{2i} = m_1 v_{1f} + m_2 v_{2f}$$

- Kinetic energy conservation:

$$ \frac{1}{2} m_1 v_{1i}^2 + \frac{1}{2} m_2 v_{2i}^2 = \frac{1}{2} m_1 v_{1f}^2 + \frac{1}{2} m_2 v_{2f}^2 $$

These equations can be solved simultaneously to find the final velocities after collision.

One-Dimensional Collisions

Most introductory problems deal with collisions along a straight line. The key steps include:

- Applying conservation of momentum.
- Applying conservation of kinetic energy (for elastic collisions).
- Solving for unknown velocities.

Example Problem:

Two billiard balls collide elastically, with masses $m_1 = m_2$, initial velocities $v_{1i} = 3\, \text{m/s}$, $v_{2i} = -2\, \text{m/s}$. Find their velocities after collision.

Solution Approach:

- Use momentum conservation to set up an equation.
- Use kinetic energy conservation (since elastic).
- Solve for v_{1f} and v_{2f}.

Two-Dimensional Collisions

Real-world collisions often involve vectors in multiple directions, requiring the resolution of velocities into components.

Key points:

- Resolve velocities into x and y components.
- Apply conservation laws separately along each axis.
- Use vector addition to find resultant velocities post-collision.

Impulse and Its Role in Collisions

Impulse is the change in momentum caused by a force applied over a specific time interval:

$$\text{Impulse} (J) = \Delta p = F \Delta t$$

In collision analysis:

- Impulse equals the change in momentum of the object.
- It helps quantify the effect of forces during impacts, especially when forces are large but act over very short times.

Real-World Applications of Momentum and Collisions

Understanding linear momentum and collisions is crucial across various fields:

- Automotive Safety: Designing crumple zones and airbags to absorb impact energy, reducing injuries.
- Sports: Analyzing how balls rebound in tennis, soccer, or billiards.
- Particle Physics: Studying collision experiments to reveal fundamental particles.
- Engineering: Designing machinery and materials to withstand impacts.

Summary and Key Takeaways

- Linear momentum is a vector quantity conserved in isolated systems.
- Collisions are classified as elastic, inelastic, or perfectly inelastic, based on energy conservation.
- Conservation of momentum is fundamental to solving collision problems.
- In elastic collisions, kinetic energy is also conserved; in inelastic collisions, some energy is dissipated.
- Impulse connects force and momentum change, especially during short-duration impacts.
- Both one-dimensional and two-dimensional collision analyses are essential for understanding real-world phenomena.

Final Thoughts

Mastering Chapter 9: Linear Momentum and Collisions provides a crucial foundation in classical mechanics. By understanding how objects interact through collisions and how momentum is conserved, students and professionals can analyze complex systems, predict outcomes, and design safer, more efficient technologies. The principles covered here are not just academic?they are actively shaping innovations across disciplines, from sports to space exploration.

Whether you're tackling homework problems, preparing for exams, or applying these concepts in practical scenarios, a solid grasp of linear momentum and collision principles unlocks a deeper understanding of the dynamic world around us.

QuestionAnswer What is the principle of conservation of linear momentum in collisions? The principle states that in an isolated system with no external forces, the total linear momentum before a collision is equal to the total linear momentum after the collision. How do elastic and inelastic collisions differ in terms of energy conservation? In elastic collisions, both kinetic energy and momentum are conserved, whereas in inelastic collisions, only momentum is conserved and some kinetic energy is transformed into other forms of energy like heat or deformation. What is the significance of the coefficient of restitution in collision analysis? The coefficient of restitution measures the 'bounciness' of a collision, indicating how much kinetic energy is conserved; it ranges from 0 (perfectly inelastic) to 1 (perfectly elastic). How is the concept of impulse related to linear momentum in collisions? Impulse is the change in momentum experienced by an object when a force is applied over a period of time, mathematically expressed as impulse = force × time = change in momentum. Can you explain the concept of center of mass in the context of collisions? The center of mass is the point where the mass of a system is considered to be concentrated; during collisions, the motion of the center of mass remains unaffected by internal forces, moving uniformly if no external forces act on the system.

Related keywords: linear momentum, collisions, elastic collision, inelastic collision, conservation of momentum, impulse, collision theory, momentum formula, collision analysis, impact force

Igcse physics momentum example questions

igcse physics momentum example questions are an essential part of exam preparation for students aiming to excel in physics. Understanding how to approach these questions can significantly improve your problem-solving skills and boost your confidence during assessments. Momentum, a fundamental concept in physics, involves the quantity of motion an object possesses and is crucial in understanding collisions, safety features, and various real-life applications. This article provides a comprehensive guide to common IGCSE physics momentum example questions, including strategies for solving them, key concepts, and practice problems to help you master this topic.

Understanding the Concept of Momentum in IGCSE Physics

Before diving into example questions, it?s important to grasp the core principles related to

momentum.

What is Momentum?

Momentum (p) is defined as the product of an object?s mass (m) and its velocity (v):

- $p = m \times v$

This quantity is a vector, meaning it has both magnitude and direction. The SI unit for momentum is kilogram meter per second ($\mathrm{kg\,m/s}$).

Conservation of Momentum

One of the most important principles in physics is the law of conservation of momentum:

- In a closed system with no external forces, the total momentum before an event (like a collision) equals the total momentum after.
- This principle helps solve many momentum-related problems involving collisions and explosions.

Common Types of Momentum Questions in IGCSE Physics

Understanding the typical question formats will help you prepare effectively.

1. Calculating Momentum

Questions often ask you to find the momentum of an object given its mass and velocity.

2. Applying Conservation of Momentum

Problems involving collisions or explosions where the initial and final momenta are used to find unknown variables.

3. Using Impulse and Force

Questions that relate change in momentum (impulse) to force and time, often involving the impulse-momentum theorem:

- $F \times \Delta t = \Delta p$

4. Analyzing Collisions and Explosions

Real-world scenarios like car crashes or ball strikes, requiring understanding of elastic and inelastic collisions.

Strategies for Solving IGCSE Physics Momentum Questions

Mastering momentum questions involves a combination of understanding physics principles and applying mathematical techniques.

Step-by-Step Approach

- Read the question carefully to identify the known and unknown variables.
- Decide which physics principle applies?most commonly, conservation of momentum.
- Write down the relevant formulae, such as $p = m \times v$ or the conservation of momentum equation.
- Substitute the known values into the formulae.\n
- Rearrange equations to solve for the unknown variable.
- Perform calculations carefully, paying attention to units and directions.
- Check your answer to ensure it makes sense physically and mathematically.

Tips for Success

- Always consider the direction of motion, as momentum is vectorial.
- Keep units consistent throughout calculations.
- Practice a variety of problems to familiarize yourself with different contexts.
- Use diagrams to visualize collisions or movements, which can help clarify the problem.

Sample IGCSE Physics Momentum Example Questions

Practicing real questions is crucial. Below are some typical example questions with detailed explanations.

Example 1: Calculating the Momentum of an Object

Question:

A 2 kg ball is rolling at a velocity of 5 m/s to the right. What is its momentum?

Solution:

Using $p = m \times v$:

- $p = 2\,kg \times 5\,m/s = 10\,kg\,m/s$

Answer: The momentum is 10 kg·m/s to the right.

Example 2: Momentum Conservation in a Collision

Question:

A 0.5 kg ball moving at 4 m/s collides with a stationary 0.5 kg ball. After the collision, the first ball moves at 1 m/s to the right, and the second moves at some unknown velocity. Find the velocity of the second ball after the collision.

Solution:

Assuming a perfectly elastic collision and using conservation of momentum:

Total initial momentum:

- $p_{initial} = (0.5\,kg \times 4\,m/s) + (0.5\,kg \times 0\,m/s) = 2\,kg\,m/s$

Total final momentum:

- $p_{final} = (0.5\,kg \times 1\,m/s) + (0.5\,kg \times v_{2})$

Set initial and final momenta equal:

- $2 = 0.5 \times 1 + 0.5 \times v_{2}$
- $2 = 0.5 + 0.5 v_{2}$
- $0.5 v_{2} = 1.5$
- $v_{2} = 3\,m/s$

Answer: The second ball moves at 3 m/s to the right after the collision.

Example 3: Impulse and Force

Question:

A 1 kg object is moving at 10 m/s. A force of 50 N is applied to it in the same direction for 2 seconds. What is its final velocity?

Solution:

Calculate change in momentum:

- $\Delta p = F \times \Delta t = 50\,N \times 2\,s = 100\,kg\,m/s$

Initial momentum:

- $p_{initial} = 1\,kg \times 10\,m/s = 10\,kg\,m/s$

Final momentum:

- $p_{final} = p_{initial} + \Delta p = 10 + 100 = 110\,kg\,m/s$

Final velocity:

- $v_{final} = p_{final} / m = 110 / 1 = 110\,m/s$

Answer: The object?s final velocity is 110 m/s in the same direction.

Practicing with Past Papers and Additional Questions

To excel in IGCSE physics momentum questions, consistent practice is key. Here are some ways to reinforce your understanding:

- Review past exam questions available in IGCSE physics practice papers.
- Work through a variety of problems, including those involving elastic and inelastic collisions.

- Create your own questions based on real-world scenarios, such as car crashes or sports collisions.
- Use online quizzes and interactive simulations to visualize momentum and collisions.

Additional Tips for Mastering Momentum in IGCSE Physics

- Focus on Vector Quantities: Always pay attention to the direction of motion; remember that momentum is a vector.
- Master the Conservation Law: Many questions rely on the conservation of momentum; ensure you understand how to apply it in different contexts.
- Pay Attention to Units: Keep units consistent and convert where necessary to avoid errors.
- Visualize the Problem: Drawing diagrams can clarify the scenario and help identify known and unknown variables.
- Review Key Formulae: Keep a list of essential formulae and principles for quick reference during exams.

Conclusion

igcse physics momentum example questions are an integral part of understanding motion and collisions in physics. By familiarizing yourself with typical question types, practicing problems regularly, and mastering the underlying principles such as conservation of momentum, you can confidently approach your exams. Remember, solving momentum questions requires a clear understanding of the concepts, careful calculations, and attention to detail. Use the strategies and example problems outlined in this guide to enhance your problem-solving skills and achieve success in your IGCSE physics assessments. Happy studying!

Understanding IGCSE Physics Momentum Example Questions: A Comprehensive Guide

When preparing for IGCSE Physics exams, one of the critical areas students need to master is the concept of momentum. Momentum questions often appear in various formats, testing your understanding of how objects move and interact. To excel, it's essential to analyze example questions thoroughly, understand the principles involved, and develop strategies for solving them efficiently. This guide aims to provide a detailed breakdown of typical IGCSE Physics momentum example questions, offering insights into approaches, common pitfalls, and step-by-step solutions to build your confidence and accuracy.

What is Momentum in Physics?

Before diving into example questions, let's briefly review the concept of momentum:

- Definition: Momentum (p) is the product of an object's mass (m) and its velocity (v):

$p = m \times v$

- Units: The SI unit of momentum is kilogram meter per second (kg·m/s).
- Principles: Momentum is a vector quantity, meaning it has both magnitude and direction.

Understanding these fundamentals helps in solving problems related to collisions, explosions, and other interactions.

Types of Momentum Questions in IGCSE Physics

IGCSE Physics momentum questions typically fall into several categories:

- Calculating momentum given mass and velocity
- Determining velocity or mass from momentum
- Analyzing collisions and explosions
- Applying the conservation of momentum
- Using the principle of impulse

By recognizing these types, students can quickly identify the key concepts needed.

Step-by-Step Approach to Solving Momentum Questions

To effectively tackle momentum questions, follow this structured approach:

- Identify what is being asked: Clarify whether you're calculating momentum, velocity, mass, or analyzing a collision.
- List known values: Write down the given data ? masses, velocities, times, etc.
- Choose the appropriate formula:

- $p = m \times v$
- Conservation of momentum: total initial momentum = total final momentum
- Impulse: change in momentum = force × time

- Apply units carefully: Ensure consistent units (e.g., kg for mass, m/s for velocity).

- Perform calculations step-by-step: Show all working to avoid errors.
- Check your answer: Verify that the units and magnitude make sense.

Common Example Questions and Solutions

Example 1: Calculating Momentum

Question:

A 0.5 kg ball is moving to the right at 4 m/s. What is its momentum?

Solution:

- Known: m = 0.5 kg, v = 4 m/s
- p = m × v = 0.5 kg × 4 m/s = 2 kg·m/s
- Answer: The momentum of the ball is 2 kg·m/s to the right.

Example 2: Finding Velocity from Momentum

Question:

A car of mass 1000 kg has a momentum of 20,000 kg·m/s. What is its velocity?

Solution:

- Known: p = 20,000 kg·m/s, m = 1000 kg
- v = p / m = 20,000 / 1000 = 20 m/s
- Answer: The car's velocity is 20 m/s in the direction of the momentum.

Example 3: Collisions and Conservation of Momentum

Question:

A 2 kg ball moving at 3 m/s collides elastically with a stationary 3 kg ball. After the collision, the 2 kg ball moves at 1 m/s in the same direction. Find the velocity of the 3 kg ball after the collision.

Solution:

- Initial momentum of system:

p_initial = (2 kg × 3 m/s) + (3 kg × 0) = 6 kg·m/s

- Final momentum:

p_final = (2 kg × 1 m/s) + (3 kg × v?)

- By conservation of momentum:

6 = 2 × 1 + 3 × v?

6 = 2 + 3v?

3v? = 4

v? = 4 / 3 ? 1.33 m/s

- Answer: The 3 kg ball moves at approximately 1.33 m/s after the collision.

Example 4: Explosive Collision (Explosion)

Question:

A 0.2 kg object moving at 5 m/s explodes into two pieces. One piece of mass 0.1 kg moves at 8 m/s to the right. Find the velocity of the other piece.

Solution:

- Initial momentum:

p_initial = 0.2 kg × 5 m/s = 1 kg·m/s to the right.

- Final momentum:

p_final = (0.1 kg × 8 m/s) + (0.1 kg × v?)

- Conservation of momentum:

1 = 0.8 + 0.1v?

0.1v? = 1 - 0.8 = 0.2

v? = 0.2 / 0.1 = 2 m/s to the right.

- Answer: The second piece moves at 2 m/s to the right.

Example 5: Impulse and Change in Momentum

Question:

A force of 50 N acts on a 0.5 kg object for 2 seconds. Find the change in momentum of the object.

Solution:

- Impulse = Force × Time = 50 N × 2 s = 100 N·s
- Change in momentum = Impulse = 100 kg·m/s
- Answer: The momentum of the object increases by 100 kg·m/s.

Tips for Mastering Momentum Questions

- Remember the vector nature: Direction matters. Be consistent with signs (+/-) depending on motion.
- Conservation of momentum: Always check if the question involves elastic or inelastic collisions to determine applicable principles.
- Units are key: Keep units consistent throughout calculations.
- Draw diagrams: Visual aids help clarify the problem, especially with multiple objects and directions.
- Practice diverse questions: Familiarity with various scenarios enhances problem-solving speed and accuracy.

Common Pitfalls and How to Avoid Them

- Mixing up units: Always convert to SI units before calculations.

- Ignoring directions: Momentum is vectorial; consider directions carefully.
- Using incorrect formulas: Ensure the formula matches the question context (e.g., conservation of momentum applies only in isolated systems).
- Forgetting to consider all objects: In collisions, account for all involved masses and velocities.

Final Thoughts

Mastering IGCSE Physics momentum example questions requires understanding the fundamental principles and practicing a variety of problems. By analyzing example questions systematically, applying the conservation of momentum correctly, and paying attention to units and directions, students can build confidence and improve their problem-solving skills. Regular practice with these types of questions will help you excel in your exams and deepen your understanding of the physics of motion.

Good luck with your studies!

QuestionAnswer What is the formula for calculating momentum in IGCSE Physics? The formula for momentum (p) is p = mass × velocity, where mass is in kilograms and velocity in meters per second. An object with a mass of 3 kg is moving at 4 m/s. What is its momentum? Its momentum is p = 3 kg × 4 m/s = 12 kg·m/s. In a collision, how does the law of conservation of momentum apply? The law states that the total momentum before the collision equals the total momentum after, assuming no external forces act on the system. A 2 kg ball moving at 5 m/s collides with a stationary 3 kg ball. If they stick together after collision, what is their combined velocity? Using conservation of momentum: (2 kg × 5 m/s) + (3 kg × 0) = (2 kg + 3 kg) × v. So, 10 = 5 × v, giving v = 2 m/s. How does increasing the mass of an object affect its momentum if velocity remains constant? Increasing the mass increases the momentum proportionally, since p = m × v. Describe what happens to the momentum of a car during a sudden stop. The car's momentum decreases rapidly to zero as the vehicle comes to rest, with the change in momentum causing forces on the brakes and passengers. Why is momentum considered a vector quantity in physics? Because it has both magnitude and direction, like velocity, and the direction affects how momentum is conserved during collisions. What is an example of an impulse in the context of momentum? An impulse occurs when a force is applied over a period of time, such as hitting a ball with a bat, which changes the ball's momentum. How can you increase the change in momentum during an impact? By increasing the force applied or the duration over which the force acts, according to the impulse-momentum theorem.

Related keywords: IGCSE physics momentum questions, momentum problem examples, physics momentum practice, IGCSE momentum exercises, momentum conservation questions, collision and explosion questions, momentum calculations IGCSE, physics momentum revision, IGCSE momentum worksheet, momentum example problems

Read the full article online: igcse physics momentum example questions